

Diagnosing Heating and Energy Transfer in Collisionless Kinetic Plasmas

J. M. TenBarge and J. Juno

Collaborators

Lev Arzamasskiy, Ammar Hakim, Greg Howes, Kris Klein, Matt Kunz

March 9, 2018




PRINCETON
UNIVERSITY

Funded by AGS-1622306 and AGS-1338944


Particle energization in turbulence


- Nonlinear cascade of MHD Alfvén waves transitions to a cascade of kinetic Alfvén waves at the ion Larmor radius.
- Most dissipation begins at ion kinetic scales and includes (but is not limited to):
 - Wave-particle interactions (Landau, transit-time, cyclotron, ...).
 - Current sheets also reconnect at ion scales and may be responsible for dissipation.
- Can we identify the energization mechanism?



Particle energization in turbulence

- Nonlinear cascade of **Censored** transitions to a cascade of **Censored** at the ion Larmor radius.
- Most dissipation begins at ion kinetic scales and includes (but is not limited to):
 - Wave-particle interactions (Landau, transit-time, cyclotron, ...).
 - Current sheets also reconnect at ion scales and may be responsible for dissipation.
- Can we identify the energization mechanism?



- 
- Introduce the field particle correlation
 - Langmuir wave damping
 - Sod shock
 - Turbulence: Driven gyrokinetic and hybrid kinetic
 - Introduce PiD
 - Langmuir wave damping
 - Sod shock

Simulation codes

<i>Gkeyll</i> [Juno et al 2018] Continuum Vlasov-Maxwell	<i>Pegasus</i> [Kunz et al 2014] Hybrid particle-in-cell	<i>AstroGK</i> [Numata et al 2010] Continuum gyrokinetics
Fully kinetic Eulerian ions and electrons	Kinetic Lagrangian ions and massless, isothermal fluid electrons	Gyrokinetic Eulerian ions and electrons — $\omega \ll \Omega_{cs}$, $k_{ } \ll k_{\perp}$, & $\delta f \ll f_0$
Up to 3x3v ions and electrons	Up to 3x3v ions and 3x electrons	Up to 3x2v ions and electrons
Non-relativistic	Non-relativistic	Non-relativistic
	Quasineutral	Quasineutral
		Cyclotron waves and entire fast/whistler mode branch ordered out
Collisionless	Collisionless ions, hyper-resistivity added to electrons	Includes Landau collision operator
		Background quantities not evolved

Field Particle Correlations

Vlasov-Poisson [Howes, Klein, & Li (2017)]

$$\frac{\partial f_s}{\partial t} + v \frac{\partial f_s}{\partial x} - \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_s}{\partial v} = 0$$

$$f_s(x, v, t) = f_{s0}(v) + \delta f_s(x, v, t)$$

Separation useful in some cases but not necessary

$$\frac{\partial \delta f_s}{\partial t} = -v \frac{\partial \delta f_s}{\partial x} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_{s0}}{\partial v} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial \delta f_s}{\partial v}$$

Multiply by $mv^2/2$ and integrate to obtain the energy equation

$$\frac{\partial W_s}{\partial t} = \int dx \int dv \frac{1}{2} m_s v^2 \left[-v \frac{\partial \delta f_s}{\partial x} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_{s0}}{\partial v} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial \delta f_s}{\partial v} \right]$$

$$\frac{\partial W_s}{\partial t} = \int dx \int dv \frac{1}{2} m_s v^2 \frac{\partial f_s}{\partial t}$$

Vlasov-Poisson [Howes, Klein, & Li (2017)]

$$\frac{\partial f_s}{\partial t} + v \frac{\partial f_s}{\partial x} - \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_s}{\partial v} = 0$$

$$f_s(x, v, t) = f_{s0}(v) + \delta f_s(x, v, t)$$

$$\frac{\partial \delta f_s}{\partial t} = -v \frac{\partial \delta f_s}{\partial x} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_{s0}}{\partial v} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial \delta f_s}{\partial v}$$

$$\frac{\partial W_s}{\partial t} = \int dx \int dv \frac{1}{2} m_s v^2 \left[-v \frac{\partial \delta f_s}{\partial x} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_{s0}}{\partial v} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial \delta f_s}{\partial v} \right]$$

$$\frac{\partial W_s}{\partial t} = \int dx \int dv \frac{1}{2} m_s v^2 \frac{\partial f_s}{\partial t}$$

In terms of fluid moments

$$\frac{1}{2} \frac{\partial P}{\partial t} = -\nabla \cdot \frac{\mathbf{Q}}{2} + qn\mathbf{u} \cdot \mathbf{E}$$

Vlasov-Poisson [Howes, Klein, & Li (2017)]

$$\frac{\partial f_s}{\partial t} + v \frac{\partial f_s}{\partial x} - \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_s}{\partial v} = 0$$

$$f_s(x, v, t) = f_{s0}(v) + \delta f_s(x, v, t)$$

$$\frac{\partial \delta f_s}{\partial t} = -v \frac{\partial \delta f_s}{\partial x} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_{s0}}{\partial v} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial \delta f_s}{\partial v}$$

$$\frac{\partial W_s}{\partial t} = \int dx \int dv \frac{1}{2} m_s v^2 \left[-v \frac{\partial \delta f_s}{\partial x} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial f_{s0}}{\partial v} + \frac{q_s}{m_s} \frac{\partial \phi}{\partial x} \frac{\partial \delta f_s}{\partial v} \right]$$

$$\frac{\partial W_s}{\partial t} = - \int dx \int dv q_s \frac{v^2}{2} \frac{\partial \delta f_s(x, v, t)}{\partial v} E(x, t) = \int dx \int dv q_s v \delta f_s(x, v, t) E(x, t)$$

$$\frac{\partial W_s}{\partial t} = - \int dx \frac{\partial \phi}{\partial x} \int dv q_s v \delta f_s = \int dx j_s E$$

Vlasov-Poisson [Howes, Klein, & Li (2017)]

$$\frac{\partial W_s}{\partial t} = - \int dx \int dv q_s \frac{v^2}{2} \frac{\partial \delta f_s(x, v, t)}{\partial v} E(x, t) = \int dx \int dv q_s v \delta f_s(x, v, t) E(x, t)$$

The field particle correlation: $C_1(v, t, \tau) = C_\tau \left(-q_s \frac{v^2}{2} \frac{\partial \delta f_s(x_0, v, t)}{\partial v}, E(x_0, t) \right)$

Alternative for cases in
which df/dv is difficult to
compute

$$C_2(v, t, \tau) = C_\tau (q_s v \delta f_s(x_0, v, t), E(x_0, t))$$

In discrete form

$$C_1(v, t_i, \tau) = \frac{1}{N} \sum_{j=i}^{i+N} q_s \frac{v^2}{2} \frac{\partial \delta f_{sj}(v)}{\partial v} E_j$$

Note that f or δf can be
used

$$t_j \equiv t(j\Delta t)$$

$$\tau = N\Delta t,$$

Vlasov-Poisson [Howes, Klein, & Li (2017)]

$$\frac{\partial W_s}{\partial t} = - \int dx \int dv q_s \frac{v^2}{2} \frac{\partial \delta f_s(x, v, t)}{\partial v} E(x, t) = \int dx \int dv q_s v \delta f_s(x, v, t) E(x, t)$$

The field particle correlation: $C_1(v, t, \tau) = C_\tau \left(-q_s \frac{v^2}{2} \frac{\partial \delta f_s(x_0, v, t)}{\partial v}, E(x_0, t) \right)$

Alternative for cases in which df/dv is difficult to compute

$$C_2(v, t, \tau) = C_\tau (q_s v \delta f_s(x_0, v, t), E(x_0, t))$$

In discrete form

$$C_1(v, t_i, \tau) = \frac{1}{N} \sum_{j=i}^{i+N} q_s \frac{v^2}{2} \frac{\partial \delta f_{sj}(v)}{\partial v} E_j$$

Note that f or δf can be used

$$t_j \equiv t(j\Delta t) \quad \tau = N\Delta t,$$

Other quantities that will appear:

$$W_s = \int w_s dx$$

$$\Delta w_s = \int_0^t (\partial w / \partial t') dt'$$

$$R = \frac{\int_{v_1}^{v_2} |C(v)| dv}{\int_{-v_{max}}^{v_{max}} |C(v)| dv}$$

$$\langle C_1 \rangle = \int_0^t C_1(v, t', \tau) dt'$$


Bow shock

$140.2 R_J$

Langmuir waves

acoustic waves


Langmuir wave test





Langmuir wave setup (Gkeyll)

$$\begin{aligned}m_p/m_e &= 1836 \\T_p/T_e &= 1 \\v_{te}/c &= 1/12 \\n &= 1 + \alpha \cos(kx), \alpha = 0.1 \\k\lambda_D &= 0.5 \\L_x &= 4\pi\lambda_D, n_x = 32\text{cells} \rightarrow 96\text{nodes} \\v_{ex} &\in [-6, 6]v_{te}, n_{vx} = 128\text{cells} \rightarrow 378\text{nodes} \\v_{px} &\in [-6, 6]v_{tp}, n_{vx} = 16\text{cells} \rightarrow 48\text{nodes}\end{aligned}$$




Initializes a standing Langmuir wave that is moderately damped, $-\gamma/\omega \simeq 0.1$







Langmuir wave result (Gkeyll)



Langmuir wave result (Gkeyll)




Langmuir wave result (Gkeyll), $\tau\omega_{pe}=0$






Langmuir wave result (Gkeyll), $\tau\omega_{pe}=6.2$


Now, we consider the field particle correlation with a finite time interval, $\tau\omega_{pe}=6.2$, which is ~ 1.5 times the period of the mode of interest



Langmuir wave result (Gkeyll), $\tau\omega_{pe}=6.2$

Now, we consider the field particle correlation with a finite time interval, $\tau\omega_{pe}=6.2$, which is ~ 1.5 times the period of the mode of interest





Sod (electrostatic) shock

Sod shock setup and evolution (Gkeyll)

$$m_p/m_e = 1836, v_{teL}/c = 0.1$$

$$n_L = 1, n_R = 0.125$$




$$T_{eL} = 0.01, T_{eR} = 0.008$$

$$T_{pL} = 0.001, T_{pR} = 0.0008$$

$$L_x = 500\lambda_D, n_x = 512\text{cells} \rightarrow 1536\text{nodes}$$

$$v_{ex} \in [-8, 8]v_{teL}, n_{vx} = 256\text{cells} \rightarrow 768\text{nodes}$$

$$v_{px} \in [-16, 16]v_{tpL}, n_{vx} = 1024\text{cells} \rightarrow 3072\text{nodes}$$



Sod shock setup and energy(Gkeyll)

$$m_p/m_e = 1836, v_{teL}/c = 0.1$$

$$n_L = 1, n_R = 0.125$$



$$T_{eL} = 0.01, T_{eR} = 0.008$$

$$T_{pL} = 0.001, T_{pR} = 0.0008$$




$$L_x = 500\lambda_D, n_x = 512\text{cells} \rightarrow 1536\text{nodes}$$

$$v_{ex} \in [-8, 8]v_{teL}, n_{vx} = 256\text{cells} \rightarrow 768\text{nodes}$$



$$v_{px} \in [-16, 16]v_{tpL}, n_{vx} = 1024\text{cells} \rightarrow 3072\text{nodes}$$




Proton shock result at $x=100\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)




Proton shock result at $x=100\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)






Heat flux leads to moderate cooling





$$\int_0^t \left(\frac{1}{2} \frac{\partial P}{\partial t'} = -\nabla \cdot \frac{\mathbf{Q}}{2} + qn\mathbf{u} \cdot \mathbf{E} \right) dt'$$




Electron shock result at $x=100\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)




Electron shock result at $x=100\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)





$$\int_0^t \left(\frac{1}{2} \frac{\partial P}{\partial t'} = -\nabla \cdot \frac{\mathbf{Q}}{2} + qn\mathbf{u} \cdot \mathbf{E} \right) dt'$$




Heat flux leads to moderate cooling




Electron shock result at $x=100\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)







$$\frac{1}{2} \frac{\partial P}{\partial t} = -\nabla \cdot \frac{\mathbf{Q}}{2} + qn\mathbf{u} \cdot \mathbf{E}$$






Heat flux leads to moderate cooling




Proton shock result at $x=175\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)




Proton shock result at $x=175\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)






Heat flux (beam)
leads to significant
heating






$$\int_0^t \left(\frac{1}{2} \frac{\partial P}{\partial t'} = -\nabla \cdot \frac{\mathbf{Q}}{2} + qn\mathbf{u} \cdot \mathbf{E} \right) dt'$$




Electron shock result at $x=175\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)



Electron shock result at $x=175\lambda_e$, $\tau\omega_{pe}=386$ (Gkeyll)



$$\int_0^t \left(\frac{1}{2} \frac{\partial P}{\partial t'} = -\nabla \cdot \frac{\mathbf{Q}}{2} + qn\mathbf{u} \cdot \mathbf{E} \right) dt'$$



Heat flux leads to significant heating and contradicts FPC

Gyrokinetic turbulence

Gyrokinetic turbulence (AstroGK)


$$m_p/m_e = 32$$

$$k_\perp \rho_p \in [0.25, 5.5] \rightarrow L_{x,y} = 25 \rho_p$$

$$(n_x, n_y, n_z, n_\lambda, n_E) = (64, 64, 32, 64, 32)$$

$$\nu_p/k_\parallel v_{tp} = 2 \times 10^{-4}$$


Driven by Langevin antenna [TenBarge et al 2014] at outer scale



Gyrokinetic turbulence (AstroGK)


$$\begin{aligned}
 m_p/m_e &= 32 \\
 k_\perp \rho_p &\in [0.25, 5.5] \rightarrow L_{x,y} = 25\rho_p \\
 (n_x, n_y, n_z, n_\lambda, n_E) &= (64, 64, 32, 64, 32) \\
 \nu_p/k_\parallel v_{tp} &= 2 \times 10^{-4}
 \end{aligned}$$

Driven by Langevin antenna [TenBarge et al 2014] at outer scale




$$f_s(\mathbf{r}, \mathbf{v}, t) = F_{0s}(v) \left(1 - \frac{q_s \phi(\mathbf{r}, t)}{T_{0s}} \right) + h_s(\mathbf{R}_s, v_\perp, v_\parallel, t)$$

$$g_s(\mathbf{R}_s, v_\perp, v_\parallel) = h_s(\mathbf{R}_s, v_\perp, v_\parallel) - \frac{q_s F_{0s}}{T_{0s}} \left\langle \phi - \frac{\mathbf{v}_\perp \cdot \mathbf{A}_\perp}{c} \right\rangle_{\mathbf{R}_s}$$





GK turbulence field particle correlation (AstroGK)



$$C_{E_{\parallel}}(\mathbf{v}, t, \tau) = C \left(-q_s \frac{v_{\parallel}^2}{2} \frac{\partial f_s(\mathbf{r}_0, \mathbf{v}, t)}{\partial v_{\parallel}}, E_{\parallel}(\mathbf{r}_0, t) \right)$$

$$C_{E_{\parallel}}(v_{\parallel}) = \int dv_{\perp} C_{E_{\parallel}}(v_{\parallel}, v_{\perp})$$

GK turbulence field particle correlation (AstroGK)



$$C_{E_{\parallel}}(\mathbf{v}, t, \tau) = C \left(-q_s \frac{v_{\parallel}^2}{2} \frac{\partial f_s(\mathbf{r}_0, \mathbf{v}, t)}{\partial v_{\parallel}}, E_{\parallel}(\mathbf{r}_0, t) \right)$$

$$C_{E_{\parallel}}(v_{\parallel}) = \int dv_{\perp} C_{E_{\parallel}}(v_{\parallel}, v_{\perp})$$

FPC identifies dominant dissipation is consistent with KAW Landau damping

Hybrid kinetic turbulence

Hybrid turbulence field particle correlation (Pegasus)

$$\begin{aligned}\beta_p &= 1 \\ L_x \times L_y \times L_z &= (20\pi)^2 \times 160\pi\rho_p \\ (n_x, n_y, n_z, n_{ppc}) &= (200, 200, 1600, 512)\end{aligned}$$

$$\begin{aligned}\beta_p &= 0.3 \\ L_x \times L_y \times L_z &= (20\pi)^2 \times 120\pi\rho_p \\ (n_x, n_y, n_z, n_{ppc}) &= (200, 200, 1200, 512)\end{aligned}$$

Driven at largest scales in domain
with a finite time correlated force
satisfying $\nabla \cdot \mathbf{F} = 0$

$$C_{\parallel}(v, t, \tau) = C \left(-q_s \frac{v^2}{2} \frac{\partial f(x_0, v, t)}{\partial v_{\parallel}}, E_{\parallel}(x_0, t) \right)$$

$$C_{\perp}(v, t, \tau) = C \left(-q_s \frac{v^2}{2} \frac{\partial f(x_0, v, t)}{\partial \mathbf{v}_{\perp}}, \mathbf{E}_{\perp}(x_0, t) \right)$$

Hybrid turbulence field particle correlation (Pegasus)

$$\begin{aligned}\beta_p &= 1 \\ L_x \times L_y \times L_z &= (20\pi)^2 \times 160\pi\rho_p \\ (n_x, n_y, n_z, n_{ppc}) &= (200, 200, 1600, 512)\end{aligned}$$

$$\begin{aligned}\beta_p &= 0.3 \\ L_x \times L_y \times L_z &= (20\pi)^2 \times 120\pi\rho_p \\ (n_x, n_y, n_z, n_{ppc}) &= (200, 200, 1200, 512)\end{aligned}$$


Driven at largest scales in domain
with a finite time correlated force
satisfying $\nabla \cdot \mathbf{F} = 0$

$$Q_{\parallel} = q\mathbf{v}_{\parallel}\mathbf{E}_{\parallel}$$

$$Q_{\perp} = q\mathbf{v}_{\perp} \cdot \mathbf{E}_{\perp}$$


Hybrid turbulence field particle correlation (Pegasus)

- Signature of Landau damping of KAWs visible, but ...
- Dominantly perpendicular heating due to high frequency stochastic heating



$$Q_{\parallel} = q\mathbf{v}_{\parallel}\mathbf{E}_{\parallel}$$

$$Q_{\perp} = q\mathbf{v}_{\perp} \cdot \mathbf{E}_{\perp}$$



Discussion of strengths and weaknesses FPC

- Provides a clear signature of energy exchange for a variety of processes, including resonant wave-particle interactions
- Signatures of other processes are yet to be identified but likely exist
- Provides a local, single point, measure of the energy exchange
 - Great for spacecraft data
 - In some cases, it may neglect the dominant source of energy transfer
- The connection to total energy exchange is simply $J_s \cdot E$
- The FPC is connected to heating but indirectly through phase mixing and collisionality

Pi-D diagnostic

Introduction of the PiD diagnostic [Yang et al (2017)]

Beginning with the standard extended MHD equations

$$\partial_t \rho_\alpha + \nabla \cdot (\rho_\alpha \mathbf{u}_\alpha) = 0,$$

$$\partial_t (\rho_\alpha \mathbf{u}_\alpha) + \nabla \cdot (\rho_\alpha \mathbf{u}_\alpha \mathbf{u}_\alpha) = -\nabla \cdot \mathbf{P}_\alpha + n_\alpha q_\alpha (\mathbf{E} + \mathbf{u}_\alpha / c \times \mathbf{B}),$$

$$\partial_t \mathcal{E}_\alpha + \nabla \cdot (\mathcal{E}_\alpha \mathbf{u}_\alpha) = -\nabla \cdot (\mathbf{P}_\alpha \cdot \mathbf{u}_\alpha) - \nabla \cdot \mathbf{h}_\alpha + n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha.$$

Re-arrange to construct the:

$$\text{Flow energy: } \partial_t \mathcal{E}_\alpha^f + \nabla \cdot (\mathcal{E}_\alpha^f \mathbf{u}_\alpha) = -\nabla \cdot (\mathbf{P}_\alpha \cdot \mathbf{u}_\alpha) + (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha + n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha.$$

$$\text{Rest-frame (thermal) energy: } \partial_t \mathcal{E}_\alpha^{th} + \nabla \cdot (\mathcal{E}_\alpha^{th} \mathbf{u}_\alpha) = -(\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha - \nabla \cdot \mathbf{h}_\alpha$$

$$\text{Electromagnetic energy: } \partial_t \mathcal{E}^m + \frac{c}{4\pi} \nabla \cdot (\mathbf{E} \times \mathbf{B}) = -\mathbf{E} \cdot \mathbf{j}$$

Introduction of the PiD diagnostic [Yang et al (2017)]

$$\partial_t \mathcal{E}_\alpha^f + \nabla \cdot (\mathcal{E}_\alpha^f \mathbf{u}_\alpha) = -\nabla \cdot (\mathbf{P}_\alpha \cdot \mathbf{u}_\alpha) + (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha + n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha.$$

$$\partial_t \mathcal{E}_\alpha^{th} + \nabla \cdot (\mathcal{E}_\alpha^{th} \mathbf{u}_\alpha) = -(\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha - \nabla \cdot \mathbf{h}_\alpha$$

$$\partial_t \mathcal{E}^m + \frac{c}{4\pi} \nabla \cdot (\mathbf{E} \times \mathbf{B}) = -\mathbf{E} \cdot \mathbf{j}$$

Integrating
over space

$$\partial_t \langle \mathcal{E}_\alpha^f \rangle = \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}_\alpha^{th} \rangle = -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}^m \rangle = -\langle \mathbf{E} \cdot \mathbf{j} \rangle,$$

Introduction of the PiD diagnostic [Yang et al (2017)]

$$\partial_t \mathcal{E}_\alpha^f + \nabla \cdot (\mathcal{E}_\alpha^f \mathbf{u}_\alpha) = -\nabla \cdot (\mathbf{P}_\alpha \cdot \mathbf{u}_\alpha) + (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha + n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha.$$

$$\partial_t \mathcal{E}_\alpha^{th} + \nabla \cdot (\mathcal{E}_\alpha^{th} \mathbf{u}_\alpha) = -(\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha - \nabla \cdot \mathbf{h}_\alpha$$


$$\partial_t \langle \mathcal{E}_\alpha^f \rangle = \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}_\alpha^{th} \rangle = -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}^m \rangle = -\langle \mathbf{E} \cdot \mathbf{j} \rangle,$$


Integrating
over space

$$\partial_t \mathcal{E}^m + \frac{c}{4\pi} \nabla \cdot (\mathbf{E} \times \mathbf{B}) = -\mathbf{E} \cdot \mathbf{j}$$



Revisiting the Langmuir wave

Langmuir wave result




$$\partial_t \langle \mathcal{E}_\alpha^f \rangle = \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}_\alpha^{th} \rangle = -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}^m \rangle = -\langle \mathbf{E} \cdot \mathbf{j} \rangle,$$

Matches Pi-D model and predictions





Revisiting the Sod shock

Sod shock spatially integrated electrons

$$\partial_t \langle \mathcal{E}_\alpha^f \rangle = \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}_\alpha^{th} \rangle = -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}^m \rangle = -\langle \mathbf{E} \cdot \mathbf{j} \rangle,$$




Sod shock spatially integrated electrons


$$\partial_t \langle \mathcal{E}_\alpha^f \rangle = \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}_\alpha^{th} \rangle = -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle,$$


$$\partial_t \langle \mathcal{E}^m \rangle = -\langle \mathbf{E} \cdot \mathbf{j} \rangle,$$



Sod shock spatial dependence electrons



$$\frac{\partial E^f}{\partial t} = -\nabla \cdot (u E^f) - \nabla \cdot (P \cdot u) + P : \nabla u + nqu \cdot E$$





$$\frac{\partial E^{th}}{\partial t} = -\nabla \cdot (u E^{th}) - \nabla \cdot q - P : \nabla u$$

Sod shock spatially integrated ions


$$\partial_t \langle \mathcal{E}_\alpha^f \rangle = \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle,$$

$$\partial_t \langle \mathcal{E}_\alpha^{th} \rangle = -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle,$$


$$\partial_t \langle \mathcal{E}^m \rangle = -\langle \mathbf{E} \cdot \mathbf{j} \rangle,$$



Sod shock spatial dependence ions




$$\frac{\partial E^f}{\partial t} = -\nabla \cdot (u E^f) - \nabla \cdot (P \cdot u) + P : \nabla u + nqu \cdot E$$




$$\frac{\partial E^{th}}{\partial t} = -\nabla \cdot (u E^{th}) - \nabla \cdot q - P : \nabla u$$

Sod shock spatial dependence ions



$$\frac{\partial E^f}{\partial t} = -\nabla \cdot (uE^f) - \nabla \cdot (P \cdot u) + P : \nabla u + nqu \cdot E$$



$$\frac{\partial E^{th}}{\partial t} = -\nabla \cdot (uE^{th}) - \nabla \cdot q - P : \nabla u$$

Strengths and weaknesses of PiD

- Agrees with total heating when integrated over entire domain
- Much like the FPC, it can be deceptive locally
- Provides little direct insight into mechanism(s) responsible for energy exchange
- Electromagnetic energy can be transferred directly to thermal energy
- In a collisional system (even weakly collisional), additional terms contribute to energy exchange

Conclusions/Future Work

- The FPC provides a relatively simple diagnostic to identify the mechanism(s) responsible for exchanging energy between the fields and particles
- The nature of the diagnostic permits its use in single point spacecraft data analysis
- In some cases, the field-particle term is not the dominant local source of particle kinetic energy
- The PiD diagnostic directly correlates with the gain or loss of thermal energy in a collisionless system
- The PiD diagnostic also provides insight into the location but not the mechanism of energy exchange and it suffers similar difficulty as the FPC when used locally
- Neither diagnostic takes collisions into account to determine irreversible heating

Conclusions/Future Work

- The FPC provides a relatively simple diagnostic to identify the mechanism(s) responsible for exchanging energy between the fields and particles
- The nature of the diagnostic permits its use in single point spacecraft data analysis
- In some cases, the field-particle term is not the dominant local source of particle kinetic energy
- The PiD diagnostic directly correlates with the gain or loss of thermal energy in a collisionless system
- The PiD diagnostic also provides insight into the location but not the mechanism of energy exchange and it suffers similar difficulty as the FPC when used locally
- Neither diagnostic takes collisions into account to determine irreversible heating
- Both diagnostics show promise and provide useful but different insights